

FertilizationManager

trees

Eurofins Agro
Binnenhaven 5
NL - 6709 PD Wageningen
The Netherlands
T sampling: Klantenservice Agro: 0888761010
T customerservice: +31 (0)88 876 1010
E customerservice@eurofins-agro.com
I www.eurofins-agro.com

Example report
P.O. Box 170
6700 AD WAGENINGEN
The Netherlands

Analysis Investigation/ordernr: 700442/003709392 Date sampling: 01-11-2023 Date report: 14-11-2023

Results	Unit	Result	Target value	low	rath.low	good	rath.high	high
Chemical	Total N stock	kg N/ha	7710	4280 - 6430				
	C/N ratio		10	13 - 17				
	N-supplying capacity	kg N/ha	140	95 - 145				
	S-plant available	kg S/ha	26	20 - 30				
	Total S stock	kg S/ha	1165	935 - 1740				
	C/S ratio		66	50 - 75				
	S-supplying capacity	kg S/ha	18	20 - 30				
	P-plant available	kg P/ha	8,1	5,2 - 8,7				
	P-soil stock	kg P/ha	545	445 - 570				
	K-plant available	kg K/ha	440	205 - 320				
	K-soil stock	kg K/ha	715	500 - 830				
	Ca-plant available	kg Ca/ha	185	210 - 490				
	Ca-soil stock	kg Ca/ha	10965	10285 - 13090				
	Mg-plant available	kg Mg/ha	670	205 - 320				
	Mg-soil stock	kg Mg/ha	955	630 - 1050				
Physical	Na-plant available	kg Na/ha	87	44 - 87				
	Na-soil stock	kg Na/ha	87	76 - 127				
	Si-plant available	g Si/ha	103830	17460 - 75660				
	Fe-plant available	g Fe/ha	9950	7280 - 13100				
	Zn-plant available	g Zn/ha	760	1460 - 2180				
	Mn-plant available	g Mn/ha	6400	5820 - 9020				
	Cu-plant available	g Cu/ha	245	115 - 190				
	Co-plant available	g Co/ha	20	15 - 25				
	B-plant available	g B/ha	680	465 - 640				
	Mo-plant available	g Mo/ha	< 10	290 - 14550				
	Se-plant available	g Se/ha	12	10 - 13				
	Acidity (pH)		6,1	> 5,9				
	C-organic	%	2,63					
	Organic matter	%	4,6					
	SOC/SOM ratio		0,57	0,45 - 0,55				
	Carbonate lime	%	0,4	2,0 - 3,0				
	Clay (<2 µm)	%	29					
	Silt (2-50 µm)	%	39					
	Sand (>50 µm)	%	27					
	<16 µm	%	41					
	Clay-humus (CEC)	mmol+/kg	233	> 189				
	CEC-saturation	%	96	> 95				
	Ca-saturation	%	81	80 - 90				
	Mg-saturation	%	12	6,0 - 10				
	K-saturation	%	2,7	2,0 - 4,0				
	Na-saturation	%	0,6	1,0 - 1,5				
	H-saturation	%	< 0,1	< 1,0				
	Al-saturation	%	< 0,1	< 1,0				

trees

Results	Unit	Result	Target value	low	rath.low	good	very good		
	Soil crumbling	score	5,3	6,0 - 8,0					
	Soil slaking	score	6,2	6,0 - 8,0					
	Risk on wind erosion	score	9,0	6,0 - 8,0					
Biological	Unit	Result	Target value	low	rath.low	good	rath.high	high	
	Moisture retention cap.	mm	49						
	Microbial biomass	mg C/kg	586	230 - 690					
	Microbial activity	mg N/kg	102	53 - 89					
	Fungal/bacterial ratio		0,6	0,6 - 0,9					

Fertilisation recommendations

The result is compared with an agricultural target value and is categorized as low, rather low, good, rather high high. This is not an appreciation as meant in ISO 17025 (par. 7.8.6).

Legislation

The fertilisation recommendations aim to achieve an agronomical optimum yield and crop quality. The recommendations do not take any legal restrictions into consideration.

Recommend.

Recommend. Unit

Soil-based recommendation (for the coming 4 years)

Phosphate (P ₂ O ₅)	0	kg/ha	When recommendations are high, it is advised to split the amount during the 4 years, for instance supply half the amount biannially. The soil based recommendation is meant to level the soil stocks of phosphorus, potassium, calcium and magnesium to the required amounts.
Potassium (K ₂ O)	30	kg/ha	
Calcium (CaO)	760	kg/ha	The lime gift is based on an optimal pH of 6,2 For every tenth increase in pH a lime gift is required of 490 kg/ha.
Magnesium (MgO)	0	kg/ha	
Lime (nw)	490	kg/ha	The amount of effective organic matter needed is calculated for a 4 year period. At the organic matter balance the yearly application of organic matter is calculated.
Effective OM	14680	kg/ha	

trees

Recommend.	Crop	Culture	Recommendation	
Crop-based recommendation (annual)				
in kg/ha	Nitrogen (N)	Avenue and park trees	Next years	84
		Conifers	Next yrs strong growth	76
		Shrubs (deciduous)	1st yr, weak growth	32
		Cotinus		64
	Sulphate (SO ₃)	Avenue and park trees	Next years	0
		Conifers	Next yrs strong growth	0
		Shrubs (deciduous)	1st yr, weak growth	0
		Cotinus		0
	Phosphate (P ₂ O ₅)	Avenue and park trees	Next years	10
		Conifers	Next yrs strong growth	10
		Shrubs (deciduous)	1st yr, weak growth	10
		Cotinus		10
	Potassium (K ₂ O)	Avenue and park trees	Next years	20
		Conifers	Next yrs strong growth	20
		Shrubs (deciduous)	1st yr, weak growth	20
		Cotinus		20
	Calcium (CaO)	Avenue and park trees	Next years	70
		Conifers	Next yrs strong growth	70
		Shrubs (deciduous)	1st yr, weak growth	70
		Cotinus		70
	Magnesium (MgO)	Avenue and park trees	Next years	0
		Conifers	Next yrs strong growth	0
		Shrubs (deciduous)	1st yr, weak growth	0
		Cotinus		0
	Sodium (Na ₂ O)	Avenue and park trees	Next years	
		Conifers	Next yrs strong growth	
		Shrubs (deciduous)	1st yr, weak growth	
		Cotinus		
	Zinc (Zn)	Avenue and park trees	Next years	0
		Conifers	Next yrs strong growth	0
		Shrubs (deciduous)	1st yr, weak growth	0
		Cotinus		0
	Manganese (Mn)	-		See the explanation.
	Copper (Cu)	Avenue and park trees	Next years	0
		Conifers	Next yrs strong growth	0
		Shrubs (deciduous)	1st yr, weak growth	0
		Cotinus		0
	Boron (B)	Avenue and park trees	Next years	0
		Conifers	Next yrs strong growth	0
		Shrubs (deciduous)	1st yr, weak growth	0
		Cotinus		0

Crop based recommendation

The crop-based recommendation is based upon the crop needs, average yields and climatic conditions and is corrected for soil nutrient stocks and the soil supplying capacity. During the growing season the SoilCropMonitor can be used for fertilization adjustments.

trees

Explanation

The results and/or the recommendations of this analysis are valid until 2027

For more information please use the following link:

<https://www.eurofins-agro.com/en/soil-analysis-explanation>

The soil based recommendation aims to maintain the soil nutrient stocks. The K, Ca and Mg recommendation will optimize the balance at the cation-exchange-capacity (CEC). It is advised to spread the soil based recommendation for nutrients and lime application across a 4 year period. When you have applied the soil based recommendation a new soil based analysis can be used to update the concentration of the nutrient stocks.

The crop based fertilization will feed the crop and improve its quality. Due to higher/lower yields and possible losses (e.g leaching) the amount of plant available nutrients can fluctuate. Therefore, we advise you to carry out a crop based analysis (Culture analysis) to determine the actual amount of available nutrients and to update the fertilization recommendations.

Look carefully at the appreciation of the nutrients on page 1. If the target values indicate that one or more nutrient quantities are very low, consult your advisor to level these quantities.

We have assumed the following yields, when calculating the crop based recommendations:

Avenue and park trees	-
Conifers	-
Shrubs (deciduous)	-
Cotinus	-

When your yield differentiates from the above, it is recommended to adjust your fertilization accordingly

Nitrogen:

The N recommendation relates to an annual dose. If possible, we recommend splitting this N dose into several applications. You can use our SoilCheck soil test in season to determine whether subsequent applications are necessary. This test measures the plant-available N (mineral N) in the soil among other things.

Sulphur:

Sulphur (S) is released by the degradation (mineralisation) of organic matter or manure. This mineralisation is performed by soil organisms. Soil organisms are not very active under colder conditions, which means not much S is released from the soil early in the spring. Therefore, it is sensible to fertilise with S for many early crops, even if the soil content is good or high.

Phosphate:

P-supplying capacity is 15 . The target range is 17 - 27
The P-buffering capacity indicates whether the P-soil stock is high enough to maintain the level of plant available P. When the buffering capacity (buffering power) is low, the plant available P will not remain on level during the growing season: it will decrease.

Potassium:**Calcium:**

Fertilization with calcium may benefit the soil structure. You can reduce the calcium soil based recommendation with the amount of calcium applied with the lime.

Manganese:

No Manganese deficiency is to be expected.

Lime:

Give the lime prior to the most lime needing crop.

Note: with liming calcium and magnesium can be added.

Soil life:

The biological soil fertility is measured by 3 characteristics, the microbial biomass, the microbial activity, and the fungal/bacterial ratio.

The acknowledgement of the measured results is based upon the amount of organic matter. There is not a recommendation given for the measured characteristics. On the basis of research projects there will be more information available.

trees

Organic matter Figure: Organic matter balance



Yearly breakdown rate (percentage) of the total organic matter content (%): 2.9

- Stock of organic matter which will remain after 1 year in the sampled layer if no (effective) organic matter is supplied.
- Total required supply of effective organic matter as a result of the degradation of the organic matter.
- Supply through crop residues (average within provided rotation scheme or crops).
- Remaining to be supplied through e.g. animal manure, green manures and/or compost.

Crop (residue)	Culture:	Input of effective organic matter
Avenue and park trees	Next years	200
Conifers	Next yrs strong growth	200
Shrubs (deciduous)	1st yr, weak growth	200
Cotinus		200
Average input/year		200

For increasing the soil organic matter content by 0.1%: 2910 kg effective organic matter per hectare is needed.

Figure: Quality of the organic matter




Organic matter consists primarily of C, N, P, S. If the organic matter contains relatively high amounts of N and/or S, this makes it attractive to soil organisms. Soil organisms happily eat this organic matter. N and S are released in the process and the amount of organic matter decreases slightly (dynamic organic matter). Organic matter can also contain a lot of C. This is generally less attractive to soil organisms (bacteria). As a result, the organic matter is not consumed as quickly by the soil organisms; making the organic matter more stable. Stable organic matter contributes - among other factors - to the workability of the soil and the looseness. Dynamic organic matter contributes primarily to the release of N and S and is therefore a source of these nutrients for the crop. The quality of the organic matter can be changed (gradually) by paying attention to the properties of soil improvers such as animal manure, compost and crop residues.

Physical

The assessment of soil structure is based on the Ca-CEC, K-CEC, and Mg-CEC ratio. Actual soil structure is - of course - not merely depending on ratio, but also on weather conditions, moisture condition of the soil, and the weight of the machinery.


Figure: Structure triangle



trees

Physical


Figure: Texture triangle



Besides clay, the silt and sand fractions are presented as well. Clay is smaller than 2 micrometer (μm), silt particles are 2-50 μm and sand particles are larger than 50 μm . The relative distribution of soil particles is used to estimate the risk of slaking. Slaking causes the soil pores to be clogged with smaller particles and degrades soil structure. The risk of slaking is greatest at 10-20% clay.

The soil crumbling - mutual binding between soil particles - is not optimal. The measures to improve soil crumbling are diverse. Considering the results, the chance of soil slaking is small.

Figure: Water retention curve



The amount of plant available water in the sampled layer is 49 mm. This is the maximum amount you should irrigate. All excess irrigation will drain off the parcel or will sink to deeper layers.

Field capacity (pF 2,0):	41,6	% moisture
Irrigation point (pF 3,3):	28,8	% moisture
Wilting point (pF 4,2):	22,0	% moisture

Crops have difficulties to obtain water when the actual moisture level is below pF 3,3. When you are able to measure the moisture level, start with irrigation if the moisture content of the parcel is at 28,8 % and irrigate 32 mm.

The actual moisture level can be measured by using a soil moisture sensor, or collect soil from ten spots in the parcel. Measure the weight of the moist soil and the weight after 24 h drying. The difference between moist and dry soil is an indication of the moisture level of the parcel.

Contact & info

Soil layer:	0 - 25 cm
Sample was taken by:	Eurofins Agro, Monsternemer
Contact sample taking:	Klantenservice Agro: 0888761010
Sampling method:	W-pattern, at least 40 sub samples, according to Eurofins Agro standard MIN 1000

If the following information is shown in the reports, this information may have been provided by the client and may affect the valuation, advice and/or analysis result:
sampling depth, crop, culture.

trees

Method		Result	Unit	Method	RvA
Results	Total nitrogen stock	2650	mg N/kg	Em: NIRS	Q
analyses	S-plant available	9,1	mg S/kg	Em: CCL3 (Gw NEN 17294-2)	
	Total sulphur stock	400	mg S/kg	Em: STT6:(Cf NEN 17294-2)	
	P-plant available	2,8	mg P/kg	Em: CCL3 (Gw NEN 15923-1)	Q
	P-soil stock	43	mg P ₂ O ₅ /100 g	Em: NIRS	
	P-soil stock	19	mg P/100 g	Em: NIRS	
	K-plant available	152	mg K/kg	Em: CCL3 (Gw NEN 17294-2)	
	K-soil stock	6,3	mmol+/kg	Em: NIRS	
	Ca-plant available	0,8	mmol Ca/l	Em: NIRS	
	Ca-soil stock	188	mmol+/kg	Em: NIRS	
	Mg-plant available	230	mg Mg/kg	Em: CCL3 (Gw NEN 17294-2)	
	Mg-soil stock	27,0	mmol+/kg	Em: NIRS	
	Na-plant available	30	mg Na/kg	Em: CCL3 (Gw NEN 17294-2)	
	Na-soil stock	1,3	mmol+/kg	Em: NIRS	
	Si-plant available	35680	µg Si/kg	Em: CCL3 (Gw NEN 17294-2)	
	Fe-plant available	3420	µg Fe/kg	Em: CCL3 (Gw NEN 17294-2)	
	Zn-plant available	260	µg Zn/kg	Em: CCL3 (Gw NEN 17294-2)	
	Mn-plant available	2200	µg Mn/kg	Em: CCL3 (Gw NEN 17294-2)	
	Cu-plant available	84	µg Cu/kg	Em: CCL3 (Gw NEN 17294-2)	Q
	Co-plant available	6,3	µg Co/kg	Em: CCL3 (Gw NEN 17294-2)	Q
	B-plant available	234	µg B/kg	Em: CCL3 (Gw NEN 17294-2)	
	Mo-plant available	< 4	µg Mo/kg	Em: CCL3 (Gw NEN 17294-2)	
	Se-plant available	4,2	µg Se/kg	Em: CCL3 (Gw NEN 17294-2)	
	Acidity (pH)	6,1		Em: NIRS	
	C-organic	2,63	%	Em: NIRS	Q
	Organic matter	4,6	%	Em: NIRS	Q
	C-inorganic	0,05	%	Em: NIRS	
	Carbonate lime	0,4	%	Em: NIRS	
	Clay (<2 µm)	29	%	Em: NIRS	
	Silt (2-50 µm)	39	%	Em: NIRS	
	Sand (>50 µm)	27	%	Em: NIRS	
	Clay-humus (CEC)	233	mmol+/kg	Em: NIRS	
	Microbial biomass	586	mg C/kg	Em: NIRS	
	Microbial activity	102	mg N/kg	Em: NIRS	
	Fungal biomass	157	mg C/kg	Em: NIRS	
	Bacterial biomass	270	mg C/kg	Em: NIRS	
	Bulk density	1164	kg/m ³	Em: NIRS	

The values stated on page 1 and 2 under 'Result' are calculated from the above mentioned analysis results.

Q Method accredited by RvA

Em: Method Eurofins Agro, Gw: Equivalent of, Cf: In conformity with

Results are reported in dry soil.

The analyses were done at Eurofins Agro, Wageningen (NL).

The results relate exclusively to the sample taken and received by Eurofins Agro, and to the material processed on 02-11-2023, and therefore to the sample analysed. For a detailed description of the sampling and analysis methods used, visit www.eurofins-agro.com